

Title	In-hospital adverse drug reactions in hospitalised older adults - a systematic review
Authors	Jennings, Emma L. M.;Murphy, Kevin D.;Gallagher, Paul F.;O'Mahony, Denis
Publication date	2018-09-17
Original Citation	Jennings, E., Murphy, K., Gallagher, P. and O'Mahony, D. (2018) '24In-Hospital Adverse Drug Reactions in Hospitalised Older Adults - A Systematic Review', 66th Annual & Scientific Meeting of the Irish Gerontological Society, Cavan, 27-29 September, in Age and Ageing, 47(suppl_5), pp. v13-v60. doi:10.1093/ageing/afy140.14
Type of publication	Conference item
Link to publisher's version	https://academic.oup.com/ageing/article/47/suppl_5/v13/5099153 - 10.1093/ageing/afy140.14
Rights	© The Author(s) 2018. Published by Oxford University Press on behalf of the British Geriatrics Society. All rights reserved.
Download date	2023-05-04 16:15:00
Item downloaded from	http://hdl.handle.net/10468/7223

In-Hospital Adverse Drug Reactions in Hospitalised Older Adults: A Systematic Review

Emma Jennings^{1,2}, Kevin Murphy³, Paul Gallagher^{1,2}, Denis O'Mahony^{1,2}

1. Department of Geriatric Medicine, Cork University Hospital, Cork, Ireland.
2. Department of Medicine, University College Cork, Cork, Ireland.
3. School of Pharmacy, University College Cork, Cork, Ireland

Introduction

- Almost ten percent of older-adults experience an adverse drug reaction [ADR] associated with acute hospitalisation
- Individual studies suggest that up to 1 in 4 experience an ADR in hospital.

This systematic review [SR] aims to evaluate in-hospital ADRs in hospitalised older-adults; frequency, culprit drug classes, severity, and clinical consequence.

Methods

This SR was conducted Following the Preferred Reporting Items for Systematic Reviews and Meta Analyses (PRISMA) (Moher et al., 2009). Two researchers (EJ, KM) screened all papers for inclusion, risk of bias and data extraction independently.

Registration

- PROSPERO registration CRD42018079095

Search Strategy

- Databases – electronic databases [PubMed, Embase and EBS-CINAHL, Cochrane Library] and library-hosted academic sources, google scholar, and grey literature.
- Search term stems – aged, ADRs, hospitalized, multi-morbid, polypharmacy and hospital-acquired [search strategy available on request from author]
- Bibliographic hand searches of relevant editorials and systematic reviews.

Paper Inclusion/Exclusion

- All languages.
- All dates up to and including the date of the final search [15/01/2018] were eligible.
- Any study that reported on ADRs either as a primary or secondary outcome in those aged 65 years or older that were hospitalised at time of ADR occurrence.
 - Review articles, systematic reviews, case reports and letters to the editor were subsequently excluded – their bibliography was hand searched for suitable studies.
- When data reported was for all ages, but there was evidence of ≥65 cohort the author was contacted requesting data for those aged 65 and over.
 - A template document for completion was provided.
 - Two attempts were made to the listed author, followed by an attempt to contact a co-author.

Study Quality / Risk of Bias Assessment

Included studies were assessed for risk of bias and study quality.


- Cochrane for randomised controlled trials
- STROBE checklist for case cohort studies
- Newcastle-Ottawa Scale for non-randomised studies

Data Extraction

We extracted data to assess percentage of study cohort ≥ 65 that experienced an ADR, reported presentation of ADRs, Drugs deemed accountable, ADR severity, ADR preventability, any measured outcomes.

Conclusion:

- There is a **lack of consistency** in methodologies used for **ADR** identification, assessment and **reporting in the literature**.
- Incident **ADRs** occurring **in-hospital** in older adults (≥ 65 years) are **common**, occurring in **1 in 5 hospitalised** older adults.
- **11** commonly prescribed **drug classes** account for **85%** of **all ADRs**.



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

Figure 1. PRISMA Flowchart outlining study selection

Results

Study selection – see PRISMA Diagram [Figure 1.]

1930 abstracts were identified, 1,779 were screened, 228 underwent full-text screening. 23 papers reporting 22 studies were included [1-23] 11 ADRs in participants ≥ 65 years, 11 ADRs all age adults (extractable data for ≥65 available in 5 studies, supplemental data provided by authors in 6 studies)

Characteristics of Included Studies

21,306 patients were included in the 22 studies; 15,769 (74%) were aged ≥65 years. 50% male, 50% female (reported in 18 studies). Polypharmacy (reported as a mean/median ≥5 medications at baseline) was reported and present in 12 studies. Multi-morbidity (reported as a mean/median number of diagnoses ≥3 at baseline) was reported in 9 and present in 8 studies.

ADR Rates

22 Studies reported ADR incidence – 2186 patients ≥ 65 years experienced ADRs in-hospital. Median 19.77% [IQR 10.44 – 25.35] Min 4.95% Max 42.19%.

Reported ADR Presentation

16 studies reported on ADR presentation (n = 13,217, 1,403 ADR presentations). 20% (283) metabolism and nutritional disorders; 17% (246) nervous system disorders; 15% (210) cardiac disorders; 13% (185) gastro-intestinal disorders; 10% (148) renal and urinary disorders; 6% (80) blood and lymphatic system disorders.

ADR Drugs

ADR associated drugs were extractable in 15 papers (1528 reported drugs in 1253 ADR patients). 85% of ADRs were associated with 11 commonly prescribed medications.

ADR Severity

14 studies reported severity (n = 13,171; reported ADRs = 1,947). 72% of reported ADRs were at least of moderate severity. 29% (560 ADRs) were severe.

ADR Preventability

5 studies assessed preventability (n = 3602, reporting 672 ADRs), 69% of reported ADRs were preventable.

Outcomes












5 papers reported on post ADR outcomes [3 length of stay [LOS], 1 LOS-death, 1 functional decline].




1 in 5 patients
≥ 65 years experiences
a clinically significant
ADR in-hospital



11 commonly
prescribed drug classes
account for 85% of ADRs

	Diuretics	22%
	Anti-thrombotics	14%
	Vitamin-K antagonists 10%, Heparin 2%, Platelet aggregation inhibitors 2%	
	Antimicrobials	12%
	Opioids	11%
	Psycholeptics	6%
	Drugs used in diabetes	4%
	Insulin 3%, oral hypoglycaemic agents 1%	
	ACE-I, ARBs	4%
	Systemic corticosteroids	3%
	Drugs for obstructive airway disease	3%
	Cardiac glycosides	3%
	Anti-hypertensives	3%

1 in 4
in-hospital ADRs
is severe


At least 2 of 3
in-hospital ADRs
are preventable

References:

1. Aydi M, Da Silva D, Martin-Bianchini J, Dal-Ripol T. Adverse Drug Reactions in Patients Hospitalized in the Intensive Care Unit of a University Hospital in Southern Brazil. *Lak Am J Pharm*. 2010 Aug;29(5):688-93.
2. Beaman L, Carlsson MC, Horvath P, Black CJ. Adverse drug reaction (ADR) occurrence and evaluation in elderly inpatients. *Pharmacoepidemiology and Drug Safety*. 1996;1(1):9-18.
3. Calderon-Ojeda C, Bustamante-Rojas C. The DOTS classification: a useful way to classify adverse drug reactions: a preliminary study in hospitalized patients. *Int J Pharm Pract*. 2010 Aug;18(4):230-5.
4. Confalonieri D, Zanetti F, Faretto F, Geronzi M, Corbelli R. Adverse drug reactions in older patients: an Italian observational prospective hospital study. *Drug Healthc Patient Saf*. 2012;4:79-80.
5. Coronado A, Truanel S, Geronzi M, Faretto F, Geronzi M, Corbelli R. Adverse drug reactions in older patients: an Italian observational prospective hospital study. *Drug Healthc Patient Saf*. 2012;4:79-80.
6. Davies C, Green CE, Taylor S, Williamson PR, Madhavan M, Pirmohamed M. Adverse drug reactions in hospital inpatients: a prospective analysis of 365 patient episodes. *PLoS one*. 2009;4(2):e4439.
7. Fernandez-Ruiz C, Gonzalez-Alvarez E, Lopez-Collado G, Alvarez-Vila A, Rodriguez-Villa A, Morin-De-la-Tor J. Hospital drug prescription and adverse drug effects in elderly patients. *Revista Clinica Espanola*. 2011 Sep;211(8):400-6.
8. Genova M, Gancheva T, Trovati J, Gancheva D, Hristakova E. A study of adverse drug reactions in hospitalized patients in relation to age. *European journal of clinical pharmacy: atención farmacéutica*. 2016;18(3):154-62.
9. Genova M, Gancheva T, Trovati J, Kiriak N, Hristakova E. Clinical relevance of drug-drug interactions in hospitalized dermatology patients. *Advances in clinical and experimental medicine: official organ Wrocław Medical University*. 2013;22(3):59-63.
10. Genova M, Gancheva T, Lazarova R, Trovati J, Kiriak N, Hristakova E. A prospective study of adverse drug reactions in a dermatology department. *Methods Find Exp Clin Pharmacol*. 2007 Mar;29(2):107-12.
11. Gonzalez-Martin G, Tanel L, Valenzuela E. Adverse drug reactions among hospitalized elderly patients. *Prospektive study*. *Revista medica de Chile*. 1997 Oct;121(10):122-36.
12. Hengert A, Pothscharer G, Samet M, Gado S, Bousvelingweiss H. Frequency and nature of adverse drug reactions in elderly inpatients of two Indian medical college hospitals. *Journal of postgraduate medicine*. 2011 Jul-Sep;57(3):189-95.
13. Lavan A, Eustice J, Doherty D, Flanagan P, Cullinan S, et al. Incident adverse drug reactions in geriatric inpatients: a multicentre observational study. *Therapeutic advances in drug safety*. 2018;9(1):1-23.
14. Lavin A, Eustice J, Doherty D, Flanagan P, Cullinan S, et al. Incident adverse drug reactions in geriatric inpatients: a multicentre observational study. *Therapeutic advances in drug safety*. 2018;9(1):1-23.
15. Mohabeh N, Shalabi G, Sakar M, Salama M, Ghazem K. Adverse drug reactions induced by cardiovascular drugs in cardiovascular care unit patients. *Pharmacoepidemiology and drug safety*. 2010;19(9):889-94.
16. Magda S, Balazs C, Kovacs A, Balazs A, Procs D, Todorovic Z. Adverse drug reactions in hospitalized cardiac patients: characteristics and risk factors. *Vigyanosmetika preprint*. 2015;7(1):137-45.
17. O'Connor KM, Gallagher P, Byrne S, O'Mahony D. Adverse drug reactions in older patients during hospitalisation: Are they predictable? Age and ageing. 2012;41(6):771-6.
18. O'Connor KM, O'Sullivan D, Gallagher P, Byrne S, O'Mahony D. Prevention of hospital-acquired adverse drug reactions in older people using screening tool of older persons' prescriptions and screening tool to alert to right treatment criteria: A cluster randomised controlled trial. *Journal of the American Geriatrics Society*. 2016;64(8):1558-66.
19. O'Connor KM, O'Sullivan D, Gallagher P, Byrne S, O'Mahony D. Prevention of hospital-acquired adverse drug reactions in older people using screening tool of older persons' prescriptions and screening tool to alert to right treatment criteria: A cluster randomised controlled trial. *Journal of the American Geriatrics Society*. 2016;64(8):1558-66.
20. O'Sullivan D, O'Mahony D, O'Connor KM, Gallagher P, Cullinan S, et al. Prevention of adverse drug reactions in hospitalised older patients using a software-supported structured pharmacist intervention: a cluster randomised controlled trial. *Drugs & ageing*. 2016;33(1):67-75.
21. REICHLE W. Complications in the care of five hundred elderly hospitalized patients. *Journal of the American Geriatrics Society*. 1985;33(11):973-81.
22. Tangiraman B, Graham-Dwyer J, Wright J, Eklund C. Adverse Drug Reactions in a Population of Hospitalized Very Elderly Patients. *Drugs & ageing*. 2012 August 01;29(8):669-79.
23. Tangiraman B, Scott G, Stevenson J, Wright J, O'Connor KM, et al. Development and validation of a risk model for predicting adverse drug reactions in older people during hospital stay. *Brighton Adverse Drug Reactions Risk (BADRR) model*. *PloS one*. 2014;9(12):e111254.